

LA-UR-14-21653

Approved for public release; distribution is unlimited.

Title: Predictions of x-ray scattering spectra in warm dense matter

Author(s): Starrett, Charles E.
Saumon, Didier
Souza, Andre N.
Perkins, David J.
Hansen, Stephanie B.

Intended for: Talk at Sandia National labs

Issued: 2014-03-12



Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Predictions of x-ray scattering spectra in warm dense matter

Charles Starrett (LANL)

Didier Saumon (LANL)

Andre Souza (U. Mich.)

David Perkins (UC, LA)

Stephanie Hansen (Sandia)

Overview

- Introduction to our model of warm dense matter
- How x-ray scattering spectra are calculated from it
- Comparisons with experiments:
 - Room temperature/pressure beryllium
 - Warm dense beryllium
 - Warm dense aluminum
- Predictions for warm dense beryllium and titanium
- Conclusions

Model for warm and hot dense matter

Electronic structure:

- e⁻ density calculated assuming superposition approximation:

$$n_e(\vec{r}) = \sum_i n_e^{PA}(|\vec{r} - \vec{R}_i|)$$

- Pseudo-atom electron density:
Calculated in spherical symmetry (inexpensive)
Average atom type approach
Kohn-Sham - DTF

Model for warm and hot dense matter

Ionic structure:


- Calculated with quantum Ornstein-Zernike relations
Exact integral equations for fluid structure
- 3 closure relations are needed → approximations:
 - Ion-ion: VMHNC
 - Electron-electron: local field corrections
 - Ion-electron: from pseudo-atom electron density.

Comparison for warm dense aluminum


Example:
Ion-ion pair distribution
functions for aluminum

Solid density (2.7g/cm^3)

Good agreement, much
quicker!



X-ray scattering



$$\hbar\omega = E_f - E_i \quad \text{Energy transfer}$$

$$k = \left| \vec{k}_f - \vec{k}_i \right| \quad \text{Momentum transfer}$$

Scattering is described by the dynamical structure factor

$$S(k, \omega)$$

Calculating x-ray spectra

- We use the Chihara formula
 - Based on chemical picture
 - Relies on clear separation between bound and screening (valence) electrons
 - Suitable for simple fluids (eg. no bonding)

$$S(k, \omega) = \left| f_I(k) + q(k) \right|^2 S_{ii}(k, \omega) \quad \leftarrow \text{elastic}$$
$$+ \bar{Z} S_{ee}(k, \omega) \quad \leftarrow \text{free-free}$$
$$+ S_{bf}(k, \omega) \quad \leftarrow \text{bound-free}$$

Calculating x-ray spectra

Elastic feature:

$$\left| f_I(k) + q(k) \right|^2 S_{ii}(k, \omega)$$

$f_I(k)$ Fourier transform of bound electron density

$q(k)$ Fourier transform of screening electron density

$$f_I(r) + q(r) = n_e^{PA}(r)$$

We already calculate this in our model!

Calculating x-ray spectra

Elastic feature:

$$|f_I(k) + q(k)|^2 S_{ii}(k, \omega)$$

$$S_{ii}(k, \omega) \approx S_{ii}(k) \delta(\omega)$$

Width of ion feature of the order \sim meV, width of x-ray probe \sim eV, so this is reasonable.

$S_{ii}(k)$ Already calculated in our model!

$S_{ii}(\omega, k)$ We plan to calculate this directly with MD

A.N. Souza *et al*, PRE (2014)

Calculating x-ray spectra

Free-free feature

$$\overline{Z} S_{ee}(k, \omega)$$

Random phase approximation (RPA)

Collisionless approximation – questionable!

Needs, free electron density, ion density, and temperature as inputs (calculated in our model)

Born-Mermin (BM)

Includes electron-ion collisions (perturbative)

Also needs ionic structure factor and interaction potential (also calculated in our model)

Calculating x-ray spectra

Bound-free feature

$$S_{bf}(k, \omega)$$

Calculate bound-free matrix elements using wave-functions from model (bound and free)

DFT wave-functions and eigenvalues – band gap problem!

W. Johnson *et al*, PRE (2012)

Validation on cold (solid) beryllium

Mattern *et al*, PRB (2012)


$$E_f = 9890 \text{ eV}$$

$$FWHM = 1.3 \text{ eV}$$


Free-free (RPA): too narrow, too strongly peaked

Bound-free: edge shifted, otherwise ok, tail in good agreement

Overall, reasonable agreement given ours is a plasma model



Warm dense beryllium: Normalization



HJ Lee *et al*, PRL (2009)

$$E_i \approx 6200 \text{ eV}$$

Mn He- α source


13 eV

$$3\rho_0 = 5.55 \text{ g/cm}^3$$

Arbitrary normalization

90° Scattered spectrum
incompatible with source!

Warm dense beryllium: Components




HJ Lee *et al*, PRL (2009)

Inelastic components are shifted to lower energies compared to probe energy

Increase left-hand peak relative to right-hand
→ Opposite to what is observed!

This implies variability in the probe beam

Warm dense beryllium: Sensitivity



HJ Lee *et al*, PRL (2009)


$\pm 30\%$ in temperature or density

Arguable that all curves fit data well, especially given source variability and arbitrary normalization

→ Large error bars on density and temperature from fit.

→ Uncertainty too large to constrain models

Warm dense aluminum



T. Ma *et al*, PRL (2013)

$$E_i \approx 17900 \text{ eV}$$

Mo $2p \rightarrow 1s$ source:
325 eV FWHM Gaussian

10 eV


$$3\rho_0 = 8.1 \text{ g/cm}^3$$

Absolute measurements

Good agreement at 69°

Strong disagreement at 111°


Warm dense aluminum



We checked:

- bound-free: compared to independent calculation (MUZE)
- Bound states against Dirac-Fock-Slater calculation
- Normalization vs. sum rules
- Exchange correlation potential
- $S_{ii}(k)$ vs. QMD
- RPA with 2 independent codes
- Validity of RPA
- Sensitivity to density and temperature ($\pm 30\%$)
- Chihara model assumptions



Warm dense aluminum



Conclusion:

→ Data at 111° cannot be reconciled with our model

Hypothetical experiment on WD beryllium



$$E_f = 9890 eV$$


$$FWHM = 1.3 eV$$

$$13 eV$$

$$3\rho_0 = 5.55 g/cm^3$$

Temperature affect:
Blurring of Fermi-
edge

Resonances in warm dense titanium



$$E_i = 4750 \text{ keV}$$

Mono-energetic source

5 eV

4.51 g/cm³

Angle: 130°

Average atom predicts resonance structures that are expected to be too sharp, due to spherical symmetry

Source spectrum would need to be narrow (~few eV) to measure width of feature

Chihara formula assumptions may break down

Next steps for theory

- Improve free-free calculation. RPA and BM questionable for WDM – could use wave functions or density of states to improve free-free
- Beyond the Chihara formula

However, the theory has not yet been proved inadequate by experiment!

Conclusions

- We have seen little sensitivity in density and temperature for experiments on Be and Al
 - May be greater sensitivity in other thermodynamic regimes
- Ability of experiments to test models hindered by source variability and large energy width of source spectrum
- Chihara formula must be used with caution – assumptions not valid at or near pressure ionization, or where binding is significant

In an ideal world...

For a probe beam width $< 5\text{eV}$ we could look at:

- Bound state \rightarrow resonance transitions
- Broadening of bound-free edge due to smearing out of bound states into bands
- Is the RPA good enough for free-free?
- Bound state energies vs. DFT energies
- Break-down of Chihara formula near pressure ionization?

In short, we could probe the microscopic physics of warm dense matter and see how well (or poorly!) the models do

Acknowledgements

- **Collaborators:**

Didier Saumon (LANL)


Andre Souza (U. Mich.)

David Andre (UC, LA)

Stephanie Hansen (Sandia)

Thanks also to T. Ma, H.J. Lee, J.D. Kress, L. Collins, G. Seidler, B. Mattern, C.J. Fontes, J. F. Benage, K. Falk and K.-U. Plagemann

Backup slides: aluminum elastic feature



Backup slides: WD beryllium elastic feature

